

RESTYLANE® LYFT FACT SHEET

Restylane Lyft is the first and only HA filler FDA-approved to restore volume and fullness to both the face and hands.^{1,2}

Treating Volume Loss, Wrinkles, and Aging Hands

Restylane Lyft contains hyaluronic acid, a natural occurring substance found in the skin that helps provide fullness and elasticity. Hyaluronic Acid diminishes as we age, causing skin to lose volume while increasing the chances for wrinkles and folds to appear. *Restylane Lyft* is formulated with Galderma's unique manufacturing process called NASHA Technology, which creates a firmer gel to provide a subtle lift to the skin.¹



Before & After

Actual patient. Treated with 0.5 mL Restylane® Lyft in each cheek, and 1 mL Restylane® Lyft in each nasolabial fold. 4 weeks after treatment. Individual results may vary.

What to expect:

- Natural-looking results and a more youthful-looking appearance¹
- Results that are clinically proven to last up to 12 months in the cheeks, and up to 6 months in facial wrinkles and hands¹
- Relatively quick procedure with little or no downtime¹
- Formulated with lidocaine which numbs the treatment area, and helps reduce pain and discomfort at the injection site¹

Restylane Lyft is one of the most versatile products in the *Restylane* portfolio and is FDA-approved to treat five areas – the **midface/cheek area**, **nasolabial folds (lines around the nose and mouth)**, **oral commissures (lines at the corner of the mouth)**, **marionette lines (vertical lines from the corners of the mouth down to the chin)**, and most recently, the **back of the hands**.¹

Formerly known as *Perlane-L*®, the product has been **available worldwide for more than 10 years**, with a **proven record of safety and efficacy**.³



Before & After

Actual patient. Treated with 1 mL Restylane® Lyft per hand. 4 weeks after treatment. Individual results may vary.



Restylane Lyft also has a high patient satisfaction rating: **96 percent of RealSelf reviewers rated the treatment as “Worth It.”**⁴

Based on 101 ratings over 24 months. Assessed July 1, 2019.

The most common side effects include bruising, redness, swelling, pain, headache, tenderness, itching at the injection site, and impaired hand function. To learn more about serious but rare side effects and full Important Safety Information visit RestylaneUSA.com.

For rewards and savings, visit www.AspireRewards.com to join ASPIRE Galderma Rewards. For more information about *Restylane*[®], visit www.RestylaneUSA.com.

Important Safety Information

Restylane® Lyft with Lidocaine is indicated for implantation into the deep dermis to superficial subcutis for the correction of moderate to severe facial folds and wrinkles, such as nasolabial folds, and for subcutaneous to supraperiosteal implantation for cheek augmentation and correction of age-related midface contour deficiencies in patients and over the age of 21.

Restylane® Lyft with Lidocaine is also indicated for injection into the subcutaneous plane in the dorsal hand to correct volume deficit in patients over the age of 21.

Restylane Lyft with Lidocaine should not be used by people with severe allergies, particularly to microorganisms known as gram-positive bacteria, or by people with serious allergies to drugs that have previously required in-hospital treatment. This product should not be used by people with bleeding disorders and should not be injected anywhere except just under the skin.

Restylane Lyft with Lidocaine should not be used in people with a known allergy to lidocaine.

The most common adverse events after initial treatment include bruising, redness, swelling, pain, headache, tenderness, itching at the injection site, and impaired hand function. Use at the site of skin sores, pimples, rashes, hives, cysts, infection, or injuries to the hand should be postponed until healing is complete. In these instances, product use could delay healing or make skin problems worse.

This product should not be injected into the blood vessels as it may cause vascular occlusion, infarction, or embolic phenomena.

Restylane Lyft with Lidocaine is available only through a licensed practitioner.

Complete Instructions for Use are available at www.RestylaneUSA.com.

© 2019 Galderma Laboratories, L.P. All trademarks are the property of their respective owners.

¹ *Restylane Lyft*. Instructions for Use. Galderma Laboratories, L.P., 2018.

² U.S. Food & Drug Administration. Medical Devices, *Restylane Lyft* with Lidocaine - P040024/S099. <https://www.fda.gov/MedicalDevices/ProductsandMedicalProcedures/DeviceApprovalsandClearances/Recently-ApprovedDevices/ucm609592.htm>. Accessed on May 18, 2018.

³ PERLANE® Injectable Gel. Summary of Safety and Effectiveness Data (PMA P040024/S6). 2007:1-16.

⁴ Real time data, Real Self website. Accessed July 1, 2019.